

X-611-68-348

PREPRINT

NASA TM X- 63331

**AN X-RAY TEST FOR SUPERDENSE  
STARS**

ELIHU BOLDT

GPO PRICE \$ \_\_\_\_\_

CSFTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 3.00Microfiche (MF) .65

SEPTEMBER 1968

ff 653 July 65



**GODDARD SPACE FLIGHT CENTER**

**GREENBELT, MARYLAND**

FACILITY FORM 602

(ACCESSION NUMBER)

N 68-34690

(THRU)


(PAGES)

5

(CODE)

1

(CATEGORY)

TMX-63331  
(NASA CR OR TMX OR AD NUMBER)

# An X-Ray Test For Superdense Stars

Elihu Boldt

NASA-Goddard Space Flight Center

Greenbelt, Maryland

Strong thermal x-ray emission from a small source ( $\sim 10^6$  cm) is a signature to be expected from a superdense star, such as a neutron star (Wheeler, 1966). It is the purpose of this letter to point out that the superdense star hypothesis for a thermal x-ray source may be tested by measuring the statistical fluctuations in the observed x-ray photon count. As an example, this is shown to be a practical test for the x-ray star Sco X-1.

As first noted by Einstein (1909-1912) and much discussed recently in connection with quantum optics (Mandel and Wolf, 1965, McLean and Pike, 1965), the photon count ( $n$ ) for a thermal source exhibits a statistical variance given by

$$(1) \quad \langle (n - \langle n \rangle)^2 \rangle = \langle n \rangle (1 + \delta)$$

where  $\delta$  is the expectation value for the photon population per unit cell ( $h^3$ ) of phase space, for each of the two spin degrees of freedom.

For independent classical particles the variance would include only the first term of (1); i.e. the variance would be  $\langle n \rangle$ . However, photons are Bosons associated with fields that interfere and hence are not completely independent. The value of  $\delta$  gives the deviation from the classical particle statistical variance in the count to be expected for thermal photons.

The quantity  $\delta$  of Equation 1 may be expressed as

$$(2) \quad \delta = \frac{\lambda^3}{2c} \cdot \left( \frac{I}{\Omega} \right)$$

where  $I$  is the spectral intensity (ergs/erg-cm<sup>2</sup>-sec) at a wavelength  $\lambda$ , from a source that subtends a solid angle  $\Omega$ ; ( $c$ ) is the velocity of light.

Note that the relation (2) between the measured variance of the photon count (1) and the measured spectral intensity ( $I$ ) at  $\lambda$  gives the solid angle  $\Omega$  subtended by the source when this information refers to thermal radiation; a small source enhances the statistical variance.

The x-ray source Sco X-1 has been measured (Gursky et al, 1965) to be a stellar (< 20") object and the optical counterpart is estimated to be at a distance of about  $8 \times 10^2$  light years (Sandage et al, 1966). If Sco X-1 is a superdense star (Shklovsky, 1967) of radius  $r \approx 10^6$  cm, then the angular size of the source is about  $10^{-9}$  arc seconds, which is far beyond the angular resolution capability ( $\sim 1$  arc second) of the grazing incidence x-ray telescope (Vaiana et al, 1968). In contrast, for the intensity from Sco X-1 measured (Grader et al, 1968) at  $\lambda = 10\text{\AA}$  ( $I \approx 10^2$  ergs/erg-cm<sup>2</sup>-sec), the situation of a superdense star of radius  $r \approx 10^6$  cm as the source of x-rays would give the significantly large value of  $\delta = 1/3$ . That this is a sensitive measure of the size of a small source may be realized by noting that for an alternate hypothetical situation where Sco X-1 is comparable in size to the sun ( $r \approx 10^{11}$  cm) the enhanced fluctuations would essentially vanish since in this case  $\delta < 10^{-10}$ .

It is a pleasure to thank William Johnston for valuable discussions.

### References

- Einstein, A.  
1909, Phys. Z. 10, 185 and 817.  
1912, "La Theorie du rayonnement et les quanta," Instituts Solway, Brussels, Conseil de Physique 1er 1911, edited by P. Langevin and L. de Broglie (Gauthier-Villars, Paris) p. 407.
- Grader, R. J., Hill, R. W., Seward, F. D., and Toor, A., 1966, Science 152, 1499.
- Gursky, H., Giacconi, R., Gorenstein, P., Waters, J. R., Oda, M., Bradt, H., Garmire, G., and Sreekantan, B. V., 1966, Astrophysical Journal 144, 1246.
- Mandel, L., and Wolf, E., 1965, Reviews of Modern Physics 37, 231.
- McLean, T. P., and Pike, E. R., 1965, Physics Letters 15, 318.
- Sandage, A. R., Osmer, P., Giacconi, R., Gorenstein, P., Gursky, H., Water, J., Bradt, H., Garmire, G., Sreekantan, B. V., Oda, M., Osawa, K., and Jugaku, J., 1966, Astrophysical Journal 146, 316.
- Shklovsky, I. S., 1967, Astrophysical Journal 148, L-1.
- Vaiana, G. S., Reidy, W. P., Zehnpfenning, T., Van Speybroeck, L., and Giacconi, R., 1968, Science 161, 564.
- Wheeler, J. A., 1966, Annual Review of Astronomy and Astrophysics 4, 393.